Module 2 : Conditional Probability

Ruben Zamar Department of Statistics UBC

January 16, 2017

Ruben Zamar Department of Statistics

- The outcome could be any element in the Sample Space, Ω .
- Sometimes the range of possibilities is restricted because of "partial information"
- Examples
 - number of shots:

partial info: we know it wasn't an "ace"

• ELEC 321 final grade:

partial info: we know it is at least a "B"

CONDITIONING EVENT

- The event *B* representing the "partial information" is called "conditioning event"
- Denote by A the event of interest
- Example (Number of Shots)

$$B = \{2, 3, ...\} = \{\text{not an "ace"}\}$$
 (conditioning event)

$$A = \{1, 3, 5, ...\} = \{\text{server wins}\}$$
 (event of interest)

• Example (Final Grade)

$$B = [70, 100] = \{ at \ least \ a \ "B" \}$$
 (conditioning event)

$$A = [80, 100] = \{an "A"\} (event of interest)$$

DEFINITION OF CONDITIONAL PROBABILITY

• Suppose that P(B) > 0

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

- The left hand side is read as "probability of A given B"
- Useful formulas:

$$P(A \cap B) = P(B) P(A|B)$$

= $P(A) P(B|A)$

CONDITIONAL PROBABILITY

- *P*(*A*|*B*), as a function of *A* (and for *B* fixed) satisfies all the probability axioms:
 - $P\left(\Omega|B\right) = P\left(\Omega \cap B\right) / P\left(B\right) = P\left(B\right) / P\left(B\right) = 1$
 - $P(A|B) \ge 0$
 - If $\{A_i\}$ are disjoint then

$$P(\cup A_i|B) = \frac{P[(\cup A_i) \cap B]}{P(B)}$$
$$= \frac{P[\cup (A_i \cap B)]}{P(B)}$$
$$= \frac{\sum P(A_i \cap B)}{P(B)} = \sum P(A_i|B)$$

EXAMPLE: NUMBER OF SHOTS

• For simplicity, suppose that points are decided in at most 8 shots, with probabilities:

Shots	1	2	3	4	5	6	7	8
Prob.	0.05	0.05	0.15	0.10	0.20	0.10	0.20	0.15

• Using the table above:

$$P(\text{Sever wins} | \text{Not an ace}) = \frac{P(\{3, 5, 7\})}{P(\{2, 3, 4, 5, 6, 7, 8\})}$$
$$= \frac{0.55}{0.95}$$
$$= 0.579$$

• Suppose that

$$P(\text{Grade is larger than } x) = \frac{100 - x}{100} = 1 - \frac{x}{100}$$

• Using the formula above:

$$P(\text{To get an "A"} | \text{To get at least a "B"}) = \frac{P([80, 100])}{P([70, 100])}$$
$$= \frac{100 - 80}{100 - 70} = \frac{20}{30}$$
$$= 0.667$$

=

- Items are submitted to a screening test before shipment
- The screening test can result in either
 - **POSITIVE** (indicating that the item may have a defect)

NEGATIVE (indicating that the item doesn't have a defect)

• Screening tests face two types of errors

FALSE POSITIVE

FALSE NEGATIVE

• For each item we have 4 possible events

Item true status:

$$D = \{\text{item is defective}\}$$

 $D^{c} = \{\text{item is not defective}\}$

Test result:

- $B = \{$ test is positive $\}$
- $B^c = \{$ test is negative $\}$

The following conditional probabilities are normally known

Sensitivity of the test: P(B|D) = 0.95 (say)

Specificity of the test: $P\left(B^{c}|D^{c}
ight)=0.99$ (say) which implies

 $P(B^{c}|D) = 0.05$ and $P(B|D^{c}) = 0.01$

• The proportion of defective items is also normally known

$$P(D) = 0.02$$
 (say)

- The following questions may be of interest:
 - What is the probability that a randomly chosen item tests positive?
 - What is the probability of defective given that the test resulted negative?
 - What is the probability of defective given that the test resulted positive?
 - What is the probability of screening error?
- We will compute these probabilities

3

(日) (周) (三) (三)

= 0.0288

- $= \ 0.02 \times 0.95 + (1 0.02) \times 0.01$
- $= P(D) P(B|D) + P(D^{c}) P(B|D^{c})$
- $P(B) = P(B \cap D) + P(B \cap D^{c})$

PROBABILITY OF TESTING POSITIVE

PROB OF DEFECTIVE GIVEN A POSITIVE TEST

$$P(D|B) = \frac{P(D \cap B)}{P(B)}$$
$$= \frac{P(D) P(B|D)}{P(B)}$$
$$= \frac{0.02 \times 0.95}{0.0288}$$
$$= 0.65972$$

э

Image: Image:

PROB OF DEFECTIVE GIVEN A NEGATIVE TEST

$$P(D|B^{c}) = \frac{P(D \cap B^{c})}{P(B^{c})}$$
$$= \frac{P(D) P(B^{c}|D)}{1 - P(B)}$$
$$= \frac{0.0098}{1 - 0.0288}$$
$$= 0.01$$

3

Image: Image:

$$P(\text{Error}) = P(D \cap B^c) + P(D^c \cap B)$$

$$= P(D) P(B^{c}|D) + P(D^{c}) P(B|D^{c})$$

$$= 0.02 \times (1 - 0.95) + (1 - 0.02) \times 0.01$$

= 0.0108

∃ ► < ∃</p>

• • • • • • • •

æ

The formula

$$P(D|B) = \frac{P(D \cap B)}{P(B)} = \frac{P(B|D)P(D)}{P(B|D)P(D) + P(B|D^{c})P(D^{c})}$$

is the simple form of Bayes' formula.

• This has been used in the "Screening Example" presented before.

• The general form of Bayes' Formula is given by

$$P(D_i|B) = \frac{P(D_i \cap B)}{P(B)} = \frac{P(B|D_i) P(D_i)}{\sum_{j=1}^k P(B|D_j) P(D_j)}$$

where D_1 , D_2 , ..., D_k is a partition of the sample space Ω :

$$\Omega = D_1 \cup D_2 \cup \cdots \cup D_k$$

$$D_i \cap D_j = \phi$$
, for $i \neq j$

- Prisoners A, B and C are to be executed
- The governor has selected one of them at random to be pardoned
- The warden knows who is pardoned, but is not allowed to tell
- Prisoner A begs the warden to let him know which one of the other two prisoners is not pardoned

- Prisoner A tells the warden: "Since I already know that one of the other two prisioners is not pardoned, you could just tell me who is that"
- Prisoner A adds: "If B is pardoned, you could give me C's name. If C is pardoned, you could give me B's name. And if I'm pardoned, you could flip a coin to decide whether to name B or C."

The warden is convinced by prisoner A's arguments and tells him: "B is not pardoned"

Result: Given the information provided by the Warden, C is now twice more likely to be pardoned than A!

Why? Check the derivations below:

NOTATION:

b =

$$A = \{A \text{ is pardoned}\}$$
$$B = \{B \text{ is pardoned}\}$$
$$C = \{C \text{ is pardoned}\}$$
$$\{\text{The warden says "B is not pardoned"}$$

Clearly

$$P(A) = P(B) = P(C) = \frac{1}{3}$$

$$P(b|B) = 0$$
 (warden never lies)
 $P(b|A) = 1/2$ (warden flips a coin)
 $P(b|C) = 1$ (warden cannot name A)

æ

→

Image: A image: A

}

By the Bayes' formula:

$$P(A|b) = \frac{P(b|A) P(A)}{P(b|A) P(A) + P(b|B) P(B) + P(b|C) P(C)}$$

= $\frac{\frac{1}{2} \times \frac{1}{3}}{\frac{1}{2} \times \frac{1}{3} + 0 \times \frac{1}{3} + 1 \times \frac{1}{3}} = \frac{1}{3}$
e
$$P(C|b) = 1 - P(A|b) = 1 - \frac{1}{2} - \frac{2}{3}$$

Hence

$$P(C|b) = 1 - P(A|b) = 1 - \frac{1}{3} = \frac{2}{3}$$

Image: Image:

æ

SCREENING EXAMPLE II

- The tested items have two components: " c_1 " and " c_2 "
- Suppose

$$D_1 \hspace{.1in} = \hspace{.1in} \{ ext{Only component "} c_1 " \hspace{.1in} ext{is defective } \} \hspace{.1in}, \hspace{.1in} P\left(D_1
ight) = 0.01$$

$$D_2 = \{ ext{Only component "} c_2 " ext{ is defective } \}, \quad P\left(D_2
ight) = 0.008$$

$$D_3 = \{ {
m Both \ components \ are \ defective } \}$$
 , $P\left(D_3
ight) = 0.002$

$$D_4 \hspace{.1in} = \hspace{.1in} \{ { t Both \hspace{.05in} components \hspace{.05in} are \hspace{.05in} non \hspace{.05in} defective \hspace{.05in} \} \hspace{.05in} , \hspace{.1in} P\left(D_4
ight) = 0.98$$

SCREENING EXAMPLE II (continued)

Let

$$B = \{$$
Screening test is positive $\}$

Suppose

 $P(B|D_1) = 0.95$ $P(B|D_2) = 0.96$ $P(B|D_3) = 0.99$ $P(B|D_4) = 0.01$

▲ロト ▲掃 ▶ ▲ 臣 ▶ ▲ 臣 ▶ ─ 臣 ─ のへで

SOME QUESTIONS OF INTEREST

- The following questions may be of interest:
 - What is the probability of testing positive?
 - What is the probability that component " c_i " (i = 1, 2) is defective when the test resulted positive?
 - What is the probability that the item is defective when the test resulted negative?
 - What is the probability both components are defective when the test resulted positive?
 - What is the probability of testing error?
- We will compute these probabilities

$P(B) = P(B \cap D_1) + P(B \cap D_2) + P(B \cap D_3) + P(B \cap D_4)$

 $= \quad 0.01 \times 0.95 + 0.008 \times 0.96 + 0.002 \times 0.99 + 0.98 \times 0.01$

= 0.02896

Notice that the probability of defective is

$$P(D) = 0.01 + 0.008 + 0.002 = 0.02$$

$$P(D_1|B) = \frac{0.01 \times 0.95}{0.02896} = 0.32804$$

$$P(D_2|B) = \frac{0.008 \times 0.96}{0.02896} = 0.26519$$

$$P(D_3|B) = \frac{0.002 \times 0.99}{0.02896} = 0.06837$$

$$P(D_4|B) = \frac{0.98 \times 0.01}{0.02896} = 0.33840$$

Ruben Zamar Department of Statistics

3

・ロト ・ 日 ト ・ ヨ ト ・ ヨ ト

$$P(B^{c}) = P(B^{c} \cap D_{1}) + P(B^{c} \cap D_{2}) + P(B^{c} \cap D_{3}) + P(B^{c} \cap D_{4})$$

 $= \quad 0.01 \times 0.05 + 0.008 \times 0.04 + 0.002 \times 0.01 + 0.98 \times 0.99$

= 0.97104

$$P(D_1|B^c) = \frac{0.01 \times 0.05}{0.97104} = 0.00051491$$

$$P(D_2|B^c) = \frac{0.008 \times 0.04}{0.97104} = 0.00032954$$

$$P(D_3|B^c) = \frac{0.002 \times 0.01}{0.97104} = 0.000020596$$

$$P(D_4|B^c) = \frac{0.98 \times 0.99}{0.97104} = 0.99913$$

$$P(\text{defective}|B^c) = 1 - 0.99913 = 0.00087$$

∃ → < ∃ →</p>

Image: A image: A

æ

• **DEFINITION:** Events A and B are independent if

$$P(A \cap B) = P(A) P(B)$$

• If A and B are independent then

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

and

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{P(A)P(B)}{P(A)} = P(B)$$

• If P(A) = 1, then A is independent of all B.

$$P(A \cap B) = P(A \cap B) + \overbrace{P(A^{c} \cap B)}^{=0} = P(B)$$
$$P(A \cap B) = \overbrace{P(A)}^{=1} P(B)$$

∃ ▶ ∢

Image: A matrix and a matrix

3

DISCUSSION (Cont)

- $\bullet\,$ Suppose that A and B are non-trivial events ($0 < P\left(A\right) < 1\,$ and $0 < P\left(B\right) < 1$)
- If A and B are mutually exclusive ($A \cap B = \phi$) then they cannot be independent because

$$P(A|B) = 0 < P(A)$$

• If $A \subset B$ then they cannot be independent because

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)}{P(B)} > P(A)$$

DISCUSSION (Cont)

- Suppose $\Omega = \{1,2,3,4,5,6,7,8,9,10\}~$ and the numbers are equally likely.
- $A = \{1, 2, 3, 4, 5\}$ and $B = \{2, 4, 6, 8\}$

•
$$P(A \cap B) = P(\{2, 4\}) = 0.20, \quad P(A) P(B) = 0.5 \times 0.4 = 0.20$$

- Hence, A and B are independent
- In terms of probabilities A is half of Ω. On the other hand A ∩ B is half of B.

• What happens if

$$P(i) = rac{i}{55}$$
 ?

•
$$P(A \cap B) = P(\{2, 4\}) = 6/55 = 0.10909$$

 $P(A) P(B) = (15/55) \times (20/55) = 0.099174$

• Hence, A and B are not independent in this case.

Definition: We say that the events $A_1, A_2, ..., A_n$ are independent if

$$P\left(A_{i_{1}}\cap A_{i_{2}}\cap\cdots\cap A_{i_{k}}\right)=P\left(A_{i_{1}}\right)P\left(A_{i_{2}}\right)\cdots P\left(A_{i_{k}}\right)$$

 $\text{for all } 1 \leq i_1 < i_2 < \cdots < i_k \leq n, \ \text{ and all } 1 \leq k \leq n.$

For example, if n = 3, then

$$P(A_{1} \cap A_{2}) = P(A_{1}) P(A_{2})$$

$$P(A_{1} \cap A_{3}) = P(A_{1}) P(A_{3})$$

$$P(A_{2} \cap A_{3}) = P(A_{2}) P(A_{3})$$

$$P(A_{1} \cap A_{2} \cap A_{3}) = P(A_{1}) P(A_{2}) P(A_{3})$$

э

SYSTEM OF INDEPENDENT COMPONENTS

In series

$$\rightarrow \quad \textbf{a} \rightarrow \textbf{b} \rightarrow \textbf{c} \rightarrow$$

• In parallel



$$A = \{ Component a works \}$$

$$B = \{\text{Component } b \text{ works}\}$$

$$C = \{\text{Component } c \text{ works}\}$$

3

・ロト ・ 日 ト ・ ヨ ト ・ ヨ ト

We assume that A, B and C are independent, that is

$$P(A \cap B \cap C) = P(A) P(B) P(C)$$

$$P(A \cap B) = P(A) P(B),$$

$$P(B \cap C) = P(B) P(C),$$

$$P(A \cap C) = P(A) P(C)$$

)

RELIABILITY CALCULATION

Problem 1: Suppose that

$$P(A) = P(B) = P(C) = 0.95.$$

Calculate the reliability of the system

$$\rightarrow \fbox{a} \rightarrow \fbox{b} \rightarrow \fbox{c} \rightarrow$$

Solution:

$$P(\text{System Works}) = P(A \cap B \cap C)$$

$$= P(A) P(B) P(C)$$

$$= 0.95^3 = 0.857$$

PRACTICE

Problem 2: Suppose that

$$P(A) = P(B) = P(C) = 0.95.$$

Calculate the reliability of the system



$$P\left(\mathsf{System works}
ight) \;\;=\;\; 1 - P\left(\mathsf{System fails}
ight)$$

$$= 1 - P(A^c \cap B^c \cap C^c)$$

$$= 1 - P(A^c) P(B^c) P(C^c)$$

$$= 1 - (1 - P(A)) (1 - P(B)) (1 - P(C))$$

Image: A matrix

$$= 1 - 0.05^3 = 0.99988$$

æ

PRACTICE

Problem 3: Suppose that

$$P(A) = P(B) = P(C) = P(D) = 0.95.$$

Calculate the reliability of the system



 $P(\text{System works}) = P(\text{subsys I works} \cap \text{subsys II works})$

= P(subsys | works) P(subsys | works)

= [1 - P(subsys I fails)] [1 - P(subsys II fails)]

$$= [1 - P(A^c \cap B^c)] [1 - P(C^c \cap D^c)]$$

$$= [1 - P(A^{c}) P(B^{c})] [1 - P(C^{c}) P(D^{c})]$$
$$= [1 - (1 - P(A)) (1 - P(B))] [1 - (1 - P(C)) (1 - P(D))]$$

$$= (1 - 0.05^2)^2 = 0.99501$$

イロト イポト イヨト イヨト 二日

Definition: We say that the events $T_1, T_2, ..., T_n$ are conditionally independent given the event B if

$$P(T_{i_1} \cap T_{i_2} \cap \cdots \cap T_{i_k} \mid B) = P(T_{i_1} \mid B) P(T_{i_2} \mid B) \cdots P(T_{i_k} \mid B)$$

for all $1 \leq i_1 < i_2 < \cdots < i_k \leq n$, and all $1 \leq k \leq n$.

For example, if
$$n = 3$$
, then
 $P(T_1 \cap T_2 \mid B) = P(T_1 \mid B) P(T_2 \mid B)$
 $P(T_1 \cap T_3 \mid B) = P(T_1 \mid B) P(T_3 \mid B)$
 $P(T_2 \cap T_3 \mid B) = P(T_2 \mid B) P(T_3 \mid B)$
 $P(T_1 \cap T_2 \cap T_3 \mid B) = P(T_1 \mid B) P(T_2 \mid B) P(T_3 \mid B)$

▲□▶ ▲圖▶ ▲圖▶ ▲圖▶ 二副 - 釣ぬ(で)

- Conditional independence doesn't imply unconditional independence and vice versa
- Conditional independence given B doesn't imply conditional independence given B^c
- However, usually both conditional independences are assumed together in applications
- We will apply this concept in Bayesian probability updating

Let S_i be the outcome of the ith test. For instance

$$egin{array}{rcl} S_1 &=& \left\{ {
m The} \ 1^{th} \ \ {
m test} \ \ {
m is \ positive}
ight\} \ S_2 &=& \left\{ {
m The} \ 2^{th} \ \ {
m test} \ \ {
m is \ negative}
ight\} \ S_3 &=& \left\{ {
m The} \ 3^{th} \ \ {
m test} \ \ {
m snegative}
ight\} \end{array}$$

and so on

The outcomes S_i (i = 1, 2, ..., n) are available in a sequential fashion.

Let
$$I_k = S_1 \cap S_2 \cap \cdots \cap S_k$$
 (data available at step k) and set

$$\pi_0 = P(E)$$

$$\pi_1 = P(E|I_1) = P(E|S_1)$$

$$\pi_2 = P(E|I_2) = P(E|S_1 \cap S_2)$$

$$\pi_3 = P(E|I_3) = P(E|S_1 \cap S_2 \cap S_3)$$

and so on

Assume that the S_i (i = 1, 2, ..., n) are independent given E and also given E^c .

Then, for k = 1, 2, ..., n

$$\pi_{k} = \frac{P(S_{k}|E)\pi_{k-1}}{P(S_{k}|E)\pi_{k-1} + P(S_{k}|E^{c})(1-\pi_{k-1})}$$

$$\pi_{k} = \frac{P(I_{k}|E) \pi_{0}}{P(I_{k}|E) \pi_{0} + P(I_{k}|E^{c}) (1 - \pi_{0})}$$

$$= \frac{P(I_{k-1} \cap S_k | E) \pi_0}{P(I_{k-1} \cap S_k | E) \pi_0 + P(I_{k-1} \cap S_k | E^c) (1 - \pi_0)}$$

$$= \frac{P(S_{k}|E) P(I_{k-1}|E) \pi_{0}}{P(S_{k}|E) P(I_{k-1}|E) \pi_{0} + P(S_{k}|E^{c}) P(I_{k-1}|E^{c}) (1-\pi_{0})}$$

э.

・ロト ・四ト ・ヨト ・ヨト

$$\pi_{k} = \frac{P(S_{k}|E) P(I_{k-1} \cap E)}{P(S_{k}|E) P(I_{k-1} \cap E) + P(S_{k}|E^{c}) P(I_{k-1} \cap E^{c})}$$

$$= \frac{P(S_{k}|E) P(I_{k-1} \cap E) / P(I_{k-1})}{\left[P(S_{k}|E) P(I_{k-1} \cap E) + P(S_{k}|E^{c}) P(I_{k-1} \cap E^{c})\right] / P(I_{k-1})}$$

$$= \frac{P(S_k|E)\pi_{k-1}}{P(S_k|E)\pi_{k-1} + P(S_k|E^c)(1-\pi_{k-1})}, \quad \pi_{k-1} = P(E|I_{k-1})$$

■ のへで

メロト メポト メヨト メヨト

Pseudo Code

• Input:

- $(S_1, S_2, S_3, ..., S_n) = (1, 0, 1, ..., 0)$ (outcomes for the *n* tests)
- $\pi = P(E)$ (prob of event of interest, for instance E= "the part is defective")

•
$$p_k = P(S_k = +|E)$$
 $k = 1, 2, ..., n$ (Sensitivity of kth test)

•
$$q_k = P(S_k = -|E^c)$$
 $k = 1, 2, ..., n$ (Specificity of kth test)

• Output
$$\pi_k = {\sf P}\left(E | S_1 \cap S_2 \cap \dots \cap S_k
ight)$$
 , $k=1,2,...,n$

Example of Input:

$$n~=~4,~~\pi=0.05$$
Test Results = $(1,1,0,1)$

k	$p_k = P\left(1 Defective ight)$	$1-q_k= P\left(1 Non \; Defective ight)$
1	$p_1 = 0.80$	$1 - q_1 = 0.05$
2	$p_2 = 0.78$	$1 - q_2 = 0.10$
3	$p_{3} = 0.85$	$1 - q_3 = 0.20$
4	<i>p</i> ₄ = 0.82	$1-q_4=0.15$

э

Image: A mathematical states and a mathem

æ

Computation of π_k

1) Initialization: Set $\pi_0 = \pi$

2) k-step:

- If $S_k = 1$, set $a = p_k$ and $b = 1 q_k$
- If $S_k=0$, set $a=1-p_k$ and $b=q_k$

3) Computing π_k :

$$\pi_k = \frac{\mathsf{a}\pi_{k-1}}{\mathsf{a}\pi_{k-1} + \mathsf{b}\left(1 - \pi_{k-1}\right)}$$

3

- When you receive an email, your spam fillter uses Bayes rule to decide whether it is spam or not.
- Basic spam filters check whether some pre-specified words appear in the email; e.g.

{diplomat,lottery,money,inheritance,president,sincerely,huge,...}.

• We consider *n* events W_i telling us whether the ith pre-specified word is in the message

Let

$$E = \{ e-mail is spam \}$$

 $W_i = \{ \text{word i is in the message} \}, i = 1, 2, ..., n$

• Assume that $W_1, W_2, ..., W_n$ are conditionally independent given E and also E^c .

- Human examination of a large number of messages is used estimate $\pi_0 = P(E)$
- The training data is also used to estimate $p_i = P(W_i|E)$ and $1 q_i = P(W_i|E^c)$
- Let $I_n = S_1 \cap S_2 \cap \cdots \cap S_n$, where S_i is either W_i or W_i^c .
- The spam filter assumes that the W_i are conditionally independent (given E and given E^c) to compute

$$P(E|I_n) = \frac{P(I_n|E) P(E)}{P(I_n|E) P(E) + P(I_n|E^c) P(E^c)}$$

Sequential Updating

• The posterior probs $\pi_k = P(E|I_k)$ (k = 1, 2, ..., n-1) can be computed sequentially using the formula

$$\pi_{k} = P(E \mid I_{k}) = \frac{P(S_{k}|E)P(E|I_{k-1})}{P(S_{k}|E)P(E|I_{k-1}) + P(S_{k}|E^{c})P(E^{c}|I_{k-1})}$$

$$= \frac{P(S_{k}|E) \pi_{k-1}}{P(S_{k}|E) \pi_{k-1} + P(S_{k}|E^{c}) (1 - \pi_{k-1})}$$

 An early decision to classify the e-mail as spam can be made if *P* (*E* | *I_k*) becomes too large (or too small).

Numerical Example

For a simple numerical example consider a case with

$$n = 8$$
 words, $P(\text{Spam}) = 0.10$

and conditional probabilities

Word	P(Word Spam)	P(Word No Spam)
W_1	0.74	0.02
W_2	0.83	0.12
W_3	0.88	0.11
W_4	0.75	0.01
W_5	0.82	0.15
W_6	0.73	0.11
W_7	0.77	0.07
W。	0.86	0.08

Word	Word Status	$P(Spam \mid I_k)$
W_1	1	0.804
W_2	0	0.443
W_3	0	0.097
W_4	1	0.889
W_5	0	0.630
W_6	1	0.919
W_7	1	0.992
W_8	1	0.999

▲□▶ ▲圖▶ ▲圖▶ ▲圖▶ 二副 - 釣ぬ(で)